

THE SYMMETRY UNDERLYING SPIN AND THE DIRAC EQUATION⁺

B.G. Sidharth*

Centre for Applicable Mathematics & Computer Sciences
B.M. Birla Science Centre, Adarsh Nagar, Hyderabad - 500063 (India)

Abstract

It is shown, in the context of a recent formulation of elementary particles in terms of, what may be called, a Quantum Mechanical Kerr-Newman metric, that spin is a consequence of a space-time cut off at the Compton wavelength and Compton time scale. On this basis, we deduce the Dirac equation from a simple coordinate transformation.

1 Introduction

The spinorial behaviour of elementary particles is a great divide which stands in the way of the unification of General Relativity and Quantum Mechanics. As noted by Wheeler and others[1, 2], the problem of introducing spin half into General Relativity, or in the reverse, introducing the curvature of space into the description of the electron had defied solution for several decades. While the concept of spin half was introduced by Uhlenbeck and Goudsmit, it is the Dirac equation which brings out the Quantum Mechanical equation of the electron in conformity with Special Relativity[3], wherein we see the emergence not only of the spin, but also the anomalous gyro magnetic ratio. As is well known the starting point for the Dirac equation is the square

⁰E-mail:birlasc@hd1.vsnl.net.in

⁺Paper at the Spin and Perturbation Theory 98, symposium, Rome, 1998.

root operator which translates itself into the well known four dimensional representation indicating the inner or extra degrees of freedom:

$$H = \sqrt{(c^2 p^2 + m^2 c^4)},$$

where

$$H = i\hbar \frac{\partial}{\partial t} \text{ and } \vec{p} = -i\hbar \vec{\nabla}$$

However, while the Dirac equation is relativistically covariant, the square root operator masks any underlying fundamental symmetry.

It is ofcourse true that one could invoke the spinorial representation of the rotation group, but the question is, why the *spinorial* representation? Our usual three dimensional space would suggest that the ordinary representation connected with the orbital angular momentum would be the natural choice. But this is not so.

In this context we will now show that if we invoke quantized space time, we are lead to the Dirac equation by a simple coordinate transformation in Minkowski space.

2 Space Time Quantization

It was suggested in the context of a Quantum Mechanical Kerr-Newman formulation of electrons that space time has in the final analysis a discrete character[4, 5], and that indeed this could be more fundamental than quantized energy[6, 7]. Infact a simple way to see this is that in the theory of the Dirac equation, the position operator is non Hermitian, that is the position coordinates are complex (cf. ref.[3]):

$$\vec{x} = c^2 \vec{p} H^{-1} + \frac{i}{2} c \hbar (\vec{\alpha} - c \vec{p} H^{-1}) H^{-1}$$

The imaginary or rapidly oscillating or Zitterbewegung part is eliminated, that is the position operator becomes Hermitian, only on averaging over space time intervals of the order of the Compton wavelength and Compton time. As we approach the Compton wavelength or time, we encounter unphysical effects, like negative energy solutions and the related Zitterbewegung. Indeed this is also true in the classical theory[8], and it was shown that by introducing the idea of a Chronon, the minimum unit of time equalling the Compton

time the difficulty could be circumvented[9]. Subsequently the concept of the chronon was investigated by others[10].

In any case such a space time cut off leads to a not only pleasing description of spin half particles (cf.ref.[4, 5]), but also to a consistent cosmology[11] which predicts an ever expanding universe as indeed has recently been confirmed by independent observations.

In a more general context, it was shown a long time ago by Snyder[12, 13] that discrete space time is entirely compatible with the Laurentz Transformation, and leads to a divergence free electrodynamics. Given a natural unit length, a , he showed that

$$\begin{aligned}[x, y] &= (ia^2/\hbar)L_z, [t, x] = (ia^2/\hbar c)M_x, \\ [y, z] &= (ia^2/\hbar)L_x, [t, y] = (ia^2/\hbar c)M_y, \\ [z, x] &= (ia^2/\hbar)L_y, [t, z] = (ia^2/\hbar c)M_z,\end{aligned}\tag{1}$$

where L_x etc. have their usual significance.

Similarly it also follows that

$$\begin{aligned}[x, p_x] &= i\hbar[1 + (a/\hbar)^2 p_x^2]; \\ [t, p_t] &= i\hbar[1 - (a/\hbar c)^2 p_t^2]; \\ [x, p_y] &= [y, p_x] = i\hbar(a/\hbar)^2 p_x p_y; \\ [x, p_t] &= c^2[p_x, t] = i\hbar(a/\hbar)^2 p_x p_t; \text{ etc.}\end{aligned}\tag{2}$$

where p_x etc. have the usual meaning.

The surprising thing about equation (1) is that the coordinates do not commute: We are automatically lead to a non commutative geometry. Equation (2) also shows a modification to the usual commutation relations. However as $a \rightarrow 0$ we recover the usual theory.

3 The Emergence of Spin

In the context of the Quantum Mechanical Kerr-Newman Black Hole formulation discussed in references[2, 4, 5] let us consider the special case where $a = \hbar/mc$, the Compton wavelength and $p_x = mc$ and so on. Substitution in equation (2) leads to

$$[x, p_x] = 2i\hbar\tag{3}$$

and similar equations.

The right hand side of (3) is double the usual value - as if the spin has been halved or the coordinate doubled. This is the typical double connectivity of spin half.

This is suggestive of the surprising fact that Quantum Mechanical spin could arise from quantized space time. This can be confirmed as follows: Let us consider a transformation of the wave function by a linear operator $U(R)$, that is

$$|\psi' \rangle = U(R)|\psi \rangle \quad (4)$$

If the transformation R is a simple infinitesimal coordinate shift in Minkowski space we will get, from (4), as is well known[14, 15]

$$\psi'(x_j) = [1 + i\epsilon(\epsilon_{ijk}x_k \frac{\partial}{\partial x_j}) + O(\epsilon^2)]\psi(x_j) \quad (5)$$

We next consider the commutation relations (1), taking a to be the Compton wavelength. One can easily verify that the choice

$$t = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \vec{x} = \begin{pmatrix} 0 & \vec{\sigma} \\ \vec{\sigma} & 0 \end{pmatrix} \quad (6)$$

provides a representation for the coordinates x and t . Equation (6) is precisely a representation of the Dirac γ matrices. Substitution of (6) in (5) now immediately leads to the Dirac equation.

Thus we have shown that given minimum space time intervals of the order of the Compton wavelength and time, a simple coordinate transformation leads to the Dirac equation. (Alternatively, we could have considered the operator $\exp(-i \epsilon^\mu p_\mu)$.)

Ofcourse the Dirac equation itself is meaningful after an averaging over these minimum space time intervals, thus completing the circle.

4 Discussion

The above conclusion should not come as a surprise. As argued in references[2, 4, 5], it is the Zitterbewegung effects within the minimum space time intervals which are symptomatic of the double connectivity of space or vice versa.

Space time *points* are classical concepts, which have no Quantum Mechanical validity, as has been pointed out by several scholars[16] - the Heisenberg Uncertainty relation forbids a physical interpretation of space time points. On the other hand it is precisely this double connectivity of spin half, which as pointed out has forbidden a reconciliation with the purely classical General Relativity. Infact a non commutative geometry as a possible solution has also been speculated upon[17].

It has also been pointed out that such a space time cut off would be a solution to the divergences encountered in Quantum Field Theory (cf.ref.[4, 13]). Indeed, we are automatically lead to a quantized electrodynamics.

Finally it may be pointed out that in the Kerr-Newman formulation referred to earlier, inertial mass, electromagnetism and gravitation naturally arise. In particular, it was argued that electromagnetism is the direct result of the opposite parity of the upper and lower (or positive and negative) components of the Dirac four spinor.

All this can be seen to be a consequence of space time quantization.

References

- [1] C.W. Misner, K.S. Thorne and J.A. Wheeler, "Gravitation", W.H. Freeman, San Francisco, 1973.
- [2] B.G. Sidharth, Gravitation and Cosmology, 4 (2) (14), 1998, p.158ff, and references therein.
- [3] P.A.M. Dirac, "Principles of Quantum Mechanics", Clarendon Press, Oxford, 1958.
- [4] B.G. Sidharth, IJPAP, **35**, 1997, p.456ff. (Also xxx.lanl.gov quant-ph 9808020).
- [5] B.G. Sidharth, IJMPA, 13(15), 1998, p.2599ff. (Also xxx.lanl.gov quant-ph 9808031).
- [6] B.G. Sidharth, "A New Approach to Locality and Causality", Vigier Symposium, Canada, 1997. (Also xxx.lanl.gov. quant-ph 9805008).

- [7] B.G. Sidharth, "Instantaneous Action at a distance in a holistic universe", Invited submission to, "Instantaneous action at a distance in Modern Physics: Pros and Contra", Eds., A. Chubykalo and R. Smirnov-Rueda, "Nova Science Books and Journals", New York (to appear).
- [8] F. Rohrlich, "Classical Charged Particles", Addison-Wesley, Reading, Mass., 1965.
- [9] P. Caldirola, "Relativity, Quanta and Cosmology", Johnson Reprint Corp., New York, 1979.
- [10] D. Finkelstein, H. Saller and Z. Tang, "Gravity, Particles and Space Time", Eds., P. Pronin, et al, World Scientific, Singapore, 1996.
- [11] B.G. Sidharth, Intl.J.Th.Phys, 37(4), 1998, p.1307ff.
- [12] H.S. Snyder, Phys. Rev. 72 (1), 1947, p.68ff.
- [13] H.S. Snyder, Phys. Rev. 71 (1), 1947, p.38ff.
- [14] S.S. Schweber, "Relativistic Quantum Field Theory", Harper and Row, New York, 1964.
- [15] J.D Bjorken and S.D Drell, "Relativistic Quantum Mechanics", McGraw-Hill Book Co., New York, 1965.
- [16] S. Sonego, Phys Letts A 208, 1995, p.1ff.
- [17] L. Smolin, in "Quantum Concepts in space and time", Eds., R. Penrose, and C.J. Isham, Clarendon Press, Oxford, 1986.